



Rhode Island Issue Brief

Influenza (Flu) Vaccination among Pregnant Women in Rhode Island: The Role of the Prenatal Care Provider

Pregnant women who were either recommended or offered influenza vaccine by their healthcare providers were nearly 6 times more likely to be vaccinated than women who were not recommended or offered the vaccine.

Background

Pregnant women have increased morbidity and mortality from seasonal influenza due to the changes that occur during pregnancy.¹⁻² Influenza vaccination with the inactivated vaccine (IIV) is the most effective way to protect pregnant women from influenza and its complications.²⁻⁴ Recent studies have demonstrated that vaccination during pregnancy significantly reduced the chance of influenza illness among infants up to six months of age.⁵⁻⁶ Vaccination of pregnant women is a key way to protect these babies who are at high risk of severe illness from influenza, but who are too young to be vaccinated.³⁻⁶

Before 2009, pregnant women had the lowest rates of seasonal influenza vaccination among all adult priority groups.^{3,7} In 2009, only about half of pregnant women received the influenza vaccine.^{3,7} Prenatal healthcare providers play a critical role in increasing rates of influenza vaccination among pregnant women by recommending and providing influenza vaccine to their patients.^{3,7-8}

Influenza Vaccination Recommendations for Pregnant Women

The Centers for Disease Control and Prevention’s (CDC) Advisory Committee on Immunization Practices (ACIP), the American College of Obstetricians and Gynecologists (ACOG) and the American Academy of Family Physicians (AAFP) recommend that women who are or plan to be pregnant during influenza season should receive the influenza vaccine as soon as possible.³⁻⁴ Inactivated influenza vaccine (IIV) is considered safe and effective during any stage of pregnancy, and is proven to benefit both the mother and baby.³⁻⁴ Live, attenuated influenza vaccine (LAIV) is not licensed for use in pregnant women.³⁻⁴

The Rhode Island Experience

Rhode Island has collected information on influenza vaccination among pregnant women through CDC’s Pregnancy Risk Assessment Monitoring System (PRAMS). PRAMS data are used to measure progress towards goals in improving the health of mothers and infants, and to identify women and infants at high risk for health problems.⁹

This Issue Brief presents the Rhode Island PRAMS data on influenza vaccination coverage rates among pregnant women in relation to the role of the prenatal care provider.

Recommended Actions for Prenatal Care Providers

- Provide a **strong** recommendation for Influenza vaccination
- Educate staff and pregnant women about the importance of influenza vaccination during pregnancy and evidence related to its safety
- Issue standing vaccine orders to prevent missed opportunities for vaccination
- Offer vaccination to pregnant women at the earliest opportunity and throughout influenza season
- Vaccinate postpartum women who were not vaccinated during pregnancy, preferably before hospital discharge or at 6 week postpartum visit
- Educate staff and postpartum women that breastfeeding is not a contraindication to vaccination
- Vaccinate all healthcare personnel in your practice to prevent spreading influenza to patients
- Establish an influenza vaccination reminder system in your practice
- Post influenza prevention announcements and provide brochures to prompt vaccination requests
- Advise family members and other close contacts of pregnant and postpartum women and infants that they should also be vaccinated against influenza

If influenza vaccine is not available in your practice, refer patients for vaccination:

- **HEALTH Information Line: 401-222-5960 or**
- <http://health.ri.gov/find/vaccinations/>

Figure 1 shows that the percentage of women who received influenza vaccination during their pregnancy increased significantly from 22% in 2004 to 74% in 2010, but it dropped in 2011 (61%). The percentage of women who reported that their healthcare provider recommended or offered influenza vaccine during their pregnancy also significantly increased from 33% in 2004 to 91% in 2010, but it dropped to 84% in 2011.




Figure 2 demonstrates that healthcare providers play a critical role in the acceptance of influenza vaccine. Pregnant women who were either recommended or offered influenza vaccine by their healthcare providers were nearly 6 times more likely to be vaccinated than women who were not recommended or offered the vaccine (69.9% vs. 11.9%).





Figure 3 reveals several reasons why women do not receive influenza vaccination during pregnancy. Vaccine safety concerns and unawareness of the importance of the vaccine during pregnancy were the major reasons for not getting vaccinated. Healthcare providers should educate women about the safety of influenza vaccine, the risk of influenza complications during pregnancy, and the protective effect of influenza vaccination on women and their infants.



References

- Dodd L, McNeil SA, Fell DB, et al. Impact of influenza exposure on rates of hospital admission and physician visits because of respiratory illness among pregnant women. *CMAJ* 2007;176(4):463-468.
- CDC. Seasonal Influenza (Flu), Pregnant Women & Influenza (Flu). Available at <http://www.cdc.gov/flu/protect/vaccine/pregnant.htm>
- CDC. Prevention and Control of Seasonal Influenza with Vaccines: Recommendations of the Advisory Committee on Immunization Practices – United States, 2013-2014. *MMWR*. September 20, 2013 / 62(RR07);1-43.
- ACOG Committee on Obstetric Practice. ACOG committee opinion number 468, 2010. Influenza vaccination during pregnancy. *Obstet Gynecol* 2010;116:1006-7.
- Zaman K, Roy E, Arifeen S, et al. Effectiveness of maternal influenza immunization in mothers and infants. *New England Journal of Medicine*. 2008;359:1555-1564.
- Benowitz I, Esposito DB, Gracey KD, Shapiro ED, Vazquez M. Influenza vaccine given to pregnant women reduces hospitalizations due to influenza in their infants. *CID* 2010; 51:1355-1361.
- Ahluwalia IB, Jamieson DJ, Rasmussen SA, et al. Correlates of Seasonal Influenza Vaccine among pregnant women in Georgia and Rhode Island. *Obstetrics & Gynecology* 2010;116:949-955.
- Rhode Island Department of Health. Influenza (Flu) Vaccination Among Pregnant Women in Rhode Island: The Role of the Prenatal Care Provider. Rhode Island Issue Brief, September 2012.
- CDC. Pregnancy Risk Assessment Monitoring System (PRAMS). Available at <http://www.cdc.gov/prams>.

For more information about Immunization, visit: www.health.ri.gov/immunization

For more information about Rhode Island Pregnancy Risk Assessment Monitoring System (PRAMS), visit: www.health.ri.gov/prams

Rhode Island PRAMS data used in this publication was made possible by a grant from the CDC.