IMMUNIZATIONS
LEAD POISONING

A REPORT ON THE STUDENTS IN YOUR LOCAL EDUCATION AGENCY

RHODE ISLAND DEPARTMENT OF HEALTH

WHAT CAN SCHOOLS AND SCHOOL DISTRICTS DO TO PROMOTE THE HEALTH AND SAFETY OF STUDENTS?

Immunization

- Ensure that school nurses complete the Rhode Island Department of Health's Annual Report of School Immunization Status for required vaccines for kindergarten and seventh grade students. This report should be completed by November 26.*
- Encourage the parents of high school students to also have their children fully immunized against vaccine-preventable diseases. High school students can either visit their primary care providers or take part in the Vaccinate Before You Graduate program.
- Encourage all students to be vaccinated against the flu every year.
- Provide annual reminders to parents about the required immunizations that children need.

More resources and information: www.health.ri.gov/immunization/for/schools
Kathy Marceau, Rhode Island Department of Health
Kathy.Marceau@health.ri.gov

Lead

- Remind parents about the importance of lead screenings for their children. Children should be screened for elevated blood-lead levels every year through six years of age.
- Advocate for safe and affordable housing in the community in which your schools are located.
- Advocate for the enforcement of housing codes and Rhode Island lead poisoning prevention regulations in the community in which your schools are located.

More resources and information: www.health.ri.gov/healthrisks/poisoning/lead
Michelle Kollett Almeida, Rhode Island Department of Health
Michelle.Kollett@health.ri.gov

* www.health.ri.gov/schoolnursesurvey

RHODE ISLAND DEPARTMENT OF HEALTH

IMMUNIZATION SURVEY RESULTS

DCYF

Kindergarten

All students entering kindergarten must have documentation of the following immunizations: five doses of Diphtheria, Tetanus, and Pertussis vaccine; three doses of Hepatitis B vaccine; two doses of Measles, Mumps, and Rubella vaccine; four doses of Polio vaccine; and two doses of Varicella (chickenpox) vaccine.*

Immunization Reporting

Local Education Agency (LEA)	Total # Schools with Grade K	# Reported	% Reported
Your LEA	0	0	0%
Rhode Island	236	233	99%

Immunization Coverage **

Local Education Agency (LEA)	Total # Students Assessed	# Fully Immunized	% Fully Immunized
Your LEA	0	0	0%
Rhode Island	11,421	10,579	93%

Number of students with no immunization record:	0
Number of students with an exemption (religious or medical):	0



Seventh Grade

All students entering seventh grade should have the following documentation of immunizations: three doses of Hepatitis B vaccine; one dose of Meningococcal conjugate (Meningitis) vaccine; two doses of Measles, Mumps, and Rubella vaccine; four doses of Polio vaccine; one dose of Tetanus, Diphtheria, and Pertussis vaccine; and two doses of Varicella (chickenpox) vaccine.*

Immunization Reporting

Local Education Agency (LEA)	Total # Schools with Grade 7	# Reported	% Reported
Your LEA	1	0	0%
Rhode Island	144	136	94%

Immunization Coverage **

Local Education Agency (LEA)	Total # Students Assessed	# Fully Immunized	% Fully Immunized
Your LEA	0	0	0%
Rhode Island	12,173	9,084	75%

Number of students with no immunization record:	0
Number of students with an exemption (religious or medical):	0



* Rhode Island Department of Health Rules and Regulations Pertaining to Immunization and Testing for Communicable Diseases (R23-1-IMM)
** Data include students enrolled in the 2013-2014 school year as of September 1, 2013. Coverage data apply only to schools that reported.

Source: Rhode Island Department of Health School Nurse Teacher Immunization Survey, 2013-2014

IMMUNIZATION SURVEY RESULTS BY SCHOOL

DCYF

Kindergarten and Seventh Grade

Immunization Coverage

School Name	Grade	Total # Students Assessed	Students Fully Immunized		# Students with No Record	# Students with an Exemption
			#	%		

Schools with No Reported Data	Grade

HEALTHY HOUSING INDICATORS

The statewide maps below show, at the census tract level, factors associated with risk for lead exposure and poor asthma control.

CHILDREN YOUNGER THAN SIX WITH FIRST-TIME ELEVATED BLOOD LEAD LEVELS, 2011-2013

Prepared by The Providence Plan, 2014
Sources: Lead Elimination Surveillance System, Rhode Island Department of Health, Rhode Island Geographic Information System
Note: Data represent the percent of children tested who had blood lead levels of 5 µg/dL or greater for the first time. Children who had elevated levels before 2011 were excluded.



CHILDREN 2 TO 17 WITH ASTHMA, 2010-2012

Prepared by The Providence Plan, 2014
Sources: Blue Cross Blue Shield of Rhode Island; Census 2010, Neighborhood Health Plan of Rhode Island, Rhode Island Department of Health, Rhode Island GIS, United Healthcare of New England
Note: Data only include children who had a doctor's office, emergency department, or inpatient hospital visit for asthma from 2010-2012. Years are not the same as for lead.



HOUSING BUILT BEFORE 1950

Prepared by The Providence Plan, 2014
Sources: US Census Bureau, 2008-2012 5-Year American Community Survey; Rhode Island Geographic Information System
Note: Data represent the percent of total housing units that was built in 1949 or earlier.



MEDIAN FAMILY INCOME, 2008-2012

Prepared by The Providence Plan, 2014
Sources: US Census Bureau, 2008-2012 5-Year American Community Survey; Rhode Island Geographic Information System
Note: Data represent the median family income in the past 12 months in 2012 inflation-adjusted dollars. Areas shown in gray are suppressed due to margins of error larger than the estimated values.



COMPOSITE MAP OF CHILDHOOD LEAD EXPOSURE, ASTHMA, MEDIAN FAMILY INCOME, AND OLDER HOUSING



Prepared by The Providence Plan, 2014

Sources: Blue Cross Blue Shield of Rhode Island; Lead Elimination Surveillance System; Neighborhood Health Plan of Rhode Island; Rhode Island Department of Education; Rhode Island Department of Health; Rhode Island Geographic Information System; United Healthcare of New England; US Census Bureau, 2008-2012 5-Year American Community Survey and 2010 Census

Method: The results of the four healthy housing indicators were standardized (using z-scores) for each census tract, averaged, and ranked into quintiles. Unreliable estimates were excluded from the ranking calculation.