

**Rocky Mountain Spotted Fever 2009-2013**  
**Rhode Island Department of Health, Division of Infectious Disease and Epidemiology**

Table 1. Frequency by Year	Year			
	2010	2011	2012	2013
Number of Cases	2	2	13	2

Table 2. Rate by Year	Year			
	2010	2011	2012	2013
Rate per 100,000	0.2	0.2	1.2	0.2

Table 3. Frequency by Agegroup and Year	Year				
	2009	2010	2011	2012	2013
0-4	0	0	0	0	0
5-9	0	0	0	0	0
10-19	0	0	0	0	0
20-29	0	1	1	3	1
30-39	0	0	0	2	0
40-49	0	0	0	3	0
50-59	0	1	0	4	1
60-69	0	0	1	1	0
70-79	0	0	0	0	0
>=80	0	0	0	0	0
Unknown	0	0	0	0	0
All	0	2	2	13	2

- Notes:**
- 1. RMSF case counts include patients classified as confirmed and probable cases.**
  - 2. Event date is generated based on the availability of data in the following order:  
 (1) illness onset date; (2) specimen collection date; (3) date of report to public health agency.**
  - 3. Rate is calculated per 100,000 population. The population denominator is based on US Census Population. Rates are based on 2008-2009 census projections and 2010 census data.**

**Rocky Mountain Spotted Fever 2009-2013**  
**Rhode Island Department of Health, Division of Infectious Disease and Epidemiology**

Table 4. Rate by Agegroup and Year	Year			
	2010	2011	2012	2013
20-29	0.7	0.7	2.0	0.7
30-39	0.0	0.0	1.6	0.0
40-49	0.0	0.0	1.9	0.0
50-59	0.7	0.0	2.6	0.7
60-69	0.0	1.0	1.0	0.0

Table 5. Frequency by Sex and Year	Year			
	2010	2011	2012	2013
Female	0	1	7	0
Male	2	1	6	2
All	2	2	13	2

Table 6. Rate by Sex and Year	Year			
	2010	2011	2012	2013
Female	0.0	0.2	1.3	0.0
Male	0.4	0.2	1.2	0.4

- Notes:**
- 1. RMSF case counts include patients classified as confirmed and probable cases.**
  - 2. Event date is generated based on the availability of data in the following order:  
 (1) illness onset date; (2) specimen collection date; (3) date of report to public health agency.**
  - 3. Rate is calculated per 100,000 population. The population denominator is based on US Census Population. Rates are based on 2008-2009 census projections and 2010 census data.**

**Rocky Mountain Spotted Fever 2009-2013**  
**Rhode Island Department of Health, Division of Infectious Disease and Epidemiology**

Table 7. Frequency by County and Year	Year			
	2010	2011	2012	2013
Kent	0	1	3	0
Newport	1	0	0	0
Providence	1	1	8	2
Washington	0	0	2	0
All	2	2	13	2

Table 8. Rate by County and Year	Year			
	2010	2011	2012	2013
Kent	0.0	0.6	1.8	0.0
Newport	1.2	0.0	0.0	0.0
Providence	0.2	0.2	1.3	0.3
Washington	0.0	0.0	1.6	0.0

Table 9. Frequency by Month and Year	Year			
	2010	2011	2012	2013
Jan	0	1	2	2
Jun	0	0	1	0
Jul	1	1	0	0
Aug	0	0	4	0
Sep	1	0	1	0
Oct	0	0	3	0

(Continued)

- Notes:**
- 1. RMSF case counts include patients classified as confirmed and probable cases.**
  - 2. Event date is generated based on the availability of data in the following order:  
 (1) illness onset date; (2) specimen collection date; (3) date of report to public health agency.**
  - 3. Rate is calculated per 100,000 population. The population denominator is based on US Census Population. Rates are based on 2008-2009 census projections and 2010 census data.**

**Rocky Mountain Spotted Fever 2009-2013**  
**Rhode Island Department of Health, Division of Infectious Disease and Epidemiology**

Table 9. Frequency by Month and Year	Year			
	2010	2011	2012	2013
Nov	0	0	2	0
All	2	2	13	2

- Notes:**
- 1. RMSF case counts include patients classified as confirmed and probable cases.**
  - 2. Event date is generated based on the availability of data in the following order:  
(1) illness onset date; (2) specimen collection date; (3) date of report to public health agency.**
  - 3. Rate is calculated per 100,000 population. The population denominator is based on US Census Population. Rates are based on 2008-2009 census projections and 2010 census data.**