

COVID-19 vaccination schedule for people with moderately to severely compromised immune systems

Vaccine	Vaccination Schedule			
Pfizer-BioNTech (age 5 to 11)	First dose	Second dose (21 days after first dose)	Additional dose (at least 28 days after second dose)	
Pfizer-BioNTech (age 12 or older)	First dose	Second dose (21 days after first dose)	Additional dose (at least 28 days after second dose)	Booster dose* (at least three months after additional dose)
Moderna (age 18 or older)	First dose	Second dose (28 days after first dose)	Additional dose (at least 28 days after second dose)	Booster dose* (at least three months after additional dose)
Johnson & Johnson (Janssen) (age 18 or older)	First dose	Additional dose (at least 28 days after first dose)	Booster dose* (at least two months after additional dose)	

People age 50 or older may get a second booster dose at least four months after first booster dose**

People age 12 or older who have compromised immune systems may get a second booster dose at least four months after their first booster dose**

People age 50 or older may get a second booster dose at least four months after first booster dose**

People age 18 or older who have compromised immune systems may get a second booster dose at least four months after their first booster dose**

*Booster doses are available and recommended to everyone age 12 or older. Any COVID-19 vaccine can be used for the booster dose in people age 18 or older, though CDC and RIDOH prefer mRNA vaccines for the primary series and booster doses.

Some groups of people may choose to get a second booster dose of **mRNA COVID-19 vaccine (Pfizer or Moderna). These groups are:

- People age 50 or older
- People age 12 or older who have moderately to severely compromised immune systems
- People age 18 or older who got Johnson & Johnson for their primary COVID-19 vaccination and Johnson & Johnson for their first booster dose

Please talk to a healthcare provider about whether a second booster dose is right for you.